

# Accuracy of Anemia Evaluation is Improved in Acutely and Chronically Ill Patients By Accounting for Volume Status

Luana Pillon MD<sup>1</sup>, Karen Mourtzikos MD<sup>2</sup>, Harold Ballard MD<sup>3</sup>, Kheng Lim MD<sup>4</sup>, Timothy Manzone MD<sup>5</sup>  
 Division of Nephrology<sup>1</sup>, Division of Nuclear Medicine<sup>2</sup>, Division of Hematology<sup>3</sup>,  
 New York University School of Medicine, New York;

Department of Radiology<sup>4</sup>, Section of Nuclear Medicine<sup>5</sup>, Christiana Care Health System, Newark, DE




## BACKGROUND

- Anemia is associated with poor prognosis in acutely and chronically ill patients – particularly in those hospitalized, with volume excess or depletion.
- While peripheral hematocrit (pHct) may provide a good estimate of red blood cell volume (RBCV) in euvoletic patients, discordance between pHct and RBCV has been reported in critical care, hematology and congestive heart failure (CHF) settings.
- Dialysis often produces large variations in pHct; 4 hours of ultrafiltration may increase pHct by > 5 % points (i.e., 19-24%).
- We considered the current anemia recommendations by KDOQI and KDIGO (goal pHct: 30-33%) and asked, **“Should volume status be considered when assessing RBCV in clinical conditions where achievement of effective perfusion and tissue oxygenation dictate management?”**
- To begin to answer this question, and to examine whether pHct is a suitable proxy for RBCV, we performed the following retrospective analysis on a large population of hospitalized patients, who had simultaneous measurements of pHct and isotope blood volume analysis (BVA), in clinical settings where volume status was in question.

## DEFINITIONS

Figure 1. Is pHct a good surrogate for RBCV?



- $TBV = PV + RBCV$  (where PV= plasma volume)
- $pHct (\% \text{ packed cell volume}) = [RBCV / (RBCV + PV)] \times 100$
- $Normalized \ Hct \ (nHct) = pHct \ (TBV / \text{Ideal TBV})$

➤ Blood Volume Analysis (BVA): Is a validated, FDA approved, tagged isotope method, whereby PV, RBCV and TBV are ascertained by a standard method and compared with established ideal values based on patient gender, height, and weight.

## HYPOTHESIS

We hypothesize that pHct is a good surrogate measure for RBCV in all states of hydration.

## METHODS

- **Design:** Retrospective study of a cohort of hospitalized patients from two large, urban, tertiary care teaching hospitals.
- **Inclusion criteria:** 627 consecutive hospitalized patients referred to nuclear medicine for tagged isotope blood volume analysis (BVA), for question of volume status (i.e. uncertain TBV, PV or RBCV). Patients included all ages (18-100 years) any race/ethnicity, gender and body mass index (BMI). Laboratory pHct was done simultaneously with BVA.
- **Exclusion criteria:** Age < 18 years, pregnancy, nursing, or iodine-allergic.
- **Study protocol:** Baseline 5mL sample of peripheral blood was drawn and sent to hospital central lab for pHct. Simultaneously, radio-isotope labeled BVA was performed using the hospital's BVA-100 (Daxor Corporation, New York, NY). 1 mL of <sup>131</sup>I-labeled albumin (<25 uCi) was injected IV over 1 min. 5mL blood samples, collected at 12, 18, 24, 30, and 36 min post injection, were assayed for radioactivity in duplicate and the results plotted (minimum three sample points, standard deviation SD <2.9%). Plasma volume (PV) was measured by extrapolating to time zero. BVA presents RBCV, PV, TBV and nHct as absolute values and as deviation (mL and %) from ideals, based on a large, healthy, heterogeneous population, stratified by gender, height and weight.
- **Statistics:** Regression Analysis. pHct and nHct were analyzed with Bland-Altman. p value of < 0.05 and 95% Confidence Intervals were considered statistically significant. Distribution of RBCV deficit and excess was categorized by deviation from norms. (SPSS, Chicago, IL).

## SUBJECTS

Table 1. n=627 Patient Characteristics		Table 2. Number (%)	
Gender	321 F, 306 M	Volume status of all patients (n=627)	
Age (years)	69.6 (19-95)	Euvoletic (TBV=Ideal TBV +/- 8%)	226 (36.0%)
BMI kg/m <sup>2</sup>	30 (18-75)	Hypovolemic (TBV>8% below Ideal TBV)	155 (24.7%)
		Hypervolemic (TBV>8% above Ideal TBV)	246 (39.2%)

## RESULTS

Table 3. Sample Patient Data

Patient n=5	Age (yrs)/sex	% Dev from ITBV (Normal +/-8%)	% Dev from IRBCV (Normal +/-10%)	% Dev from IPV (Normal +/-8%)	pHct% (goal 30-33)	nHct %	BVA Volume
1	52/M	+18	+16	+4.9	33	40.7	Hypervolemic
2	65/F	-5.8	-31.1	+7.8	32.8	32.6	Normovolemic
3	36/M	+7.4	-29.7	+28.5	33	31	Normovolemic
4	73/M	-13.3	-33.8	+7.3	31	28.6	Hypovolemic
5	81/F	-35.7	-46.5	-28.1	32	19	Hypovolemic

## RESULTS

Figure 2. Correlation between pHct (A) and nHct (B) with RBCV in hospitalized patients (n=627).




Figure 4. Correlation between pHct & deviation from ideal RBCV, by anemia status. (anemia defined as pHct<33%)




Figure 5. Correlation between nHct & deviation from ideal RBCV, by anemia status. (anemia defined as pHct<33%)




Figure 4B. Anemic patients (n=394)




Figure 5B. Anemic patients (n=394)




Figure 4C. Hypovolemic patients (n=155)




Figure 5C. Hypervolemic patients (n=246)




Figure 3. Bland-Altman analysis of pHct and nHct in hospitalized patients, by volume status.




Figure 3B. Hypovolemic patients (n=155)





Figure 3C. Hypervolemic patients (n=246)



## CONCLUSIONS

- By comparing RBCV to pHct we found that in hospitalized patients pHct is a poor surrogate for RBCV.
- pHct may underestimate RBCV in states of fluid excess (dilutional anemia) and may overestimate RBCV when PV is low (hemoconcentration).
- In states of fluid excess and depletion, pHct may be misleading if volume status is not factored into the evaluation.
- Normalized hematocrit (nHct) may provide a more reliable assessment of anemia in patients whose volume status is uncertain.

## BIBLIOGRAPHY

- Dworkin HJ, Premo M and Dees S. Comparison of red cell and whole blood volume as performed using both chromium-51-tagged red cells and iodine-125-tagged albumin and using I-131-tagged albumin and extrapolated red cell volume. Am J Med Sci 2007; 334:37-40.
- Feldschuh J and Enson Y. Prediction of the normal blood volume: Relation of blood volume to body habitus. Circulation 1977; 56:605-612.
- Androne AS, Katz SD, Lund L et al. Hemodilution is common in patients with advanced heart failure. Circulation 2003; 107:226-229.

We thank Sandra Gilbert, PhD and Shirley Rebotira, LPN, BSN for their assistance with this study.  
 Poster Session: Vascular Physiology/Renal Hemodynamics 11-6-08 (10:00 AM-12:00 PM) Poster Board Number: TH-PO250. Contact: luanapillon2002@yahoo.com